



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

JANUARY 14.

Dr. CHARLES SCHAEFFER in the chair.

Twenty-two persons present.

The death of J. Frank Knight, a member, was announced.

Pea-like Phosphorite from Polk Co., Florida.—MR. EDW. GOLD-SMITH stated that Mr. Jos. Willcox had received from Mr. E. R. Childers of Fort Meade, Fla., a specimen of rock which, it is stated, is about to be quarried for use as a fertilizer. The rock extends over a considerable area of territory that is tributary to Peace Creek, from the bottom of which stream a large amount of what is supposed to be phosphate of lime is now obtained by two companies operating there. It is essentially a phosphorite occurring in pea-like masses mixed with carbonate of lime, fragments of bones and shells, small pebbles of quartz and Limonite. Although the rock is brittle, a thin section was prepared for microscopic examination. This was accomplished by Mr. Lancaster Thomas in a way that is worth noting. The specimen was completely soaked with boiled Canada balsam prior to the grinding. The result was a transparent slide. There was observed, besides the above mentioned mixture, amorphous silica in which were imbedded acicular crystals of apatite.

The specific gravity was found to be 2.675. The rock cannot be considered rich in phosphoric acid. The latter, however, appears fairly disseminated through the whole mass. The pea-like globules were tested separately, together with fragments of bone and the dull brown-colored Limonite, phosphoric acid being found in each case. A quantitative determination of the acid has of course commercial interest but is out of the question here.

The name proposed for the rock was suggested by the pea-shaped globules. It may be considered as a variety of Phosphorite.

JANUARY 21.

The President, Dr. JOSEPH LEIDY, in the chair.

Nineteen persons present.

A paper entitled "On the Anatomy of *Aerope caffra* Fer.," by H. A. Pilsbry was presented for publication.

JANUARY 28.

Mr. J. H. REDFIELD in the chair.

Thirty persons present.

Papers under the following titles were presented for publication :—

“Remarks on *Urosalpinx perrugatus* Contr.” By Frank C. Baker.

“On the Fishes described in Müller’s Supplemental Volume to the *Systema Naturae* of Linnaeus.” By David Starr Jordan.

“Geology of Artesian Wells at Atlantic City.” By Lewis Woolman.

Mr. Carl Edelheim was elected a member.

Henry M. Stanley was elected a correspondent.

The following were ordered to be printed :—